

Review of Sudanese Medicinal Plants Screened for Antidiabetic Activity

Elbadri Elamin Osman¹, Ahmed Abdel hafiz Elshikh¹, Mohamed Ismail Garbi², Ali Mohamed Badri², Ahmed Saeed Kabbashi^{2,3*}, Bashir Elteгани³, Mubarak Siddig Hamad³, Ayat Ahmed Abdalrahman⁴, Mahmoud M. Dahab⁵ and Waleed S. Koko⁶

1. Department of Microbiology, Faculty of Pure and Applied Science, International University of Africa, Khartoum, Sudan.
2. Department of Microbiology, Faculty of Medical Laboratory Sciences, International University of Africa, Khartoum, Sudan.
3. Medicinal and Aromatic Plants and Traditional Medicine Research Institute (MAPTMRI), Khartoum, Sudan.
4. Department of Pharmacognosy, Faculty of Pharmacy, University of Medical Sciences and Technology, Khartoum, Sudan.
5. Nutritional Immunology and Functional Foods, Key Laboratory of Tropical Plant Resources, Xishuanbanna Tropical Botanical Garden, Chinese Academy of Sciences. Menglum, Mengla, Yunnan, 666303 PR China.
6. College of Science and Arts in Ar Rass, University of Qassim, K.S.A.

***Correspondence:**

Ahmed Saeed Ali Kabbashi

Email: ahmedsak88@gmail.com

Received: 23-APR-2021; Published: 01-Jun-2021;

Citation: Ahmed Saeed Ali Kabbashi, Review of Sudanese Medicinal Plants Screened for Antidiabetic Activity

Abstract

Introduction: Diabetes is a serious metabolic disorder and plenty of medical plants are used in traditional medicines to treat diabetes. These plants have no side effects, and many existing medicines are derived from the plants. The purpose of this systematic review is to study diabetes and to summarize the available treatments for this disease, focusing especially on herbal medicine.

Methods: Scientific data bases were investigated for and screening through google engine for diabetes and effective plants informatics data. However, the investigation included Science direct, PubMed, Wiley, Scopus, and Springer. Out of the 100 collected articles (published in the period between 1997 and 2018).

Results: Diabetes is mainly due to oxidative stress and an increase in reactive oxygen species that can have major effects. Many plants contain different natural antioxidants, in particular tannins, flavonoids, C and E vitamins that have the ability to maintain β -cells performance and decrease glucose levels in the blood.

Conclusion: According to published results, it can be said that medical plants are more affordable and have less side effects compared synthetic drugs and are more effective in treatment of diabetes mellitus. Also order to harness these natural resources and maximize the socioeconomic benefits derivable from Sudanese medicinal plants efforts should be geared toward research funding and deployment of Research and Development (R & D) policy framework into medicinal plants research endeavors.

Keywords: Diabetic, Medicinal plants, Sudan.

Diabetes mellitus According to WHO, the term diabetes mellitus is defined as a metabolic disorder of multiple etiology characterized by chronic hyperglycemia with disturbances of carbohydrate, fat and protein metabolism resulting from defects in insulin secretion, insulin action, or both. The effects of diabetes mellitus include long-term damage, dysfunction and failure of various organs (Mellitus et al., 2010). Diabetes mellitus may have the characteristic symptoms such as thirst, polyuria, blurred vision and loss of weight (Association, 2014).

Diabetes disease is known since antiquity, it is mentioned in the writings of the ancient civilizations, particularly Egypt, Arabia, India, China and Asia Minor (Singh, 2016). Recently, Diabetes Millets is highly prevalent worldwide when compared with many other diseases, it is fast becoming the epidemic of the 21st century (Matthews and Matthews, 2011). It is estimated that in the year 2000, there were 171 million diabetic peoples in the world. This estimation is expected to increase to reach 366 million diabetic peoples by 2030 (Ogurtsova et al., 2017). Diabetes mellitus is defined as a disease in which the body is unable to use and store glucose, properly. It is caused by the abnormality of carbohydrate metabolism which is linked to either low blood insulin level or insensitivity of target organs to insulin (Anstee et al., 2013). Moreover, there are many types of diabetes, however, the two main types of diabetes are Type 1 (insulin-dependent), where the body is completely stop producing insulin, accordingly patients must take insulin injections daily to survive (Ashcroft and Rorsman, 2012). Type 2 (non-insulin dependent) where the body is unable to produce enough insulin or the produced insulin does not work properly (Ashcroft and Rorsman, 2012). Patients of type 2 mostly have a family history of diabetes (inherited), overweight or over 40 years of age (Association, 2015). Interestingly, type 2 (non-insulin dependent) diabetes mellitus is much more prevalent than type 1 (insulin dependent) diabetes, affecting the people of both developed and developing countries (Dunstan et al., 2002). Up to 250,000 children in developing countries under the age of 14 years have type 1 diabetes; around 38,000 of these children are in Africa (Dunstan et al., 2002).

The total population in Sudan is about 34 million, with 70% of it in the Northern parts. Prevalence of type 1 diabetes is estimated to be 0.1% among school-age children 7 to 14 years old while type 2 diabetes is estimated at 10.4% among adult population (over 25 year of age) in Northern Sudan (Yagi and Yagi, 2018). Both insulin and glucagon, pancreatic endocrine hormones, are responsible for controlling blood-glucose level within the body in an adequate level based on the body needs. Normally, insulin is secreted by the β -cells found at the islets of Langerhans in response to high levels of blood sugar (Kahn et al., 2006). It potentiates the ability of muscle, red blood cells, and fat cells to absorb sugar out of the blood and consume it in other metabolic processes, which restore the sugar levels to the normal level (Singab et al., 2014).

Natural compounds may be possible alternatives for the

treatment of diabetes and its associated complications at the behest of failure in sustainable cure from the modern drugs. In view of the effectiveness, safety and low cost production of plant medicines, consumption of the source in the daily diet can potentially manage and may even reduce the risk of the disease (Patwardhan and Partwardhan, 2005). The medicinal plants have played a remarkable and characteristic role as therapeutic agents in the treatment of chronic pathologies that are linked to diabetes before the introduction of insulin in 1922 (He et al., 2015). In this connection the plant sources (if taken in the form of diet) are considered to be very effective and safe because they potentially manage and control the risk factors associated with the diabetes (Association, 2016).

Recently, some medicinal plants have been reported to be useful in diabetes worldwide and have been used empirically as antidiabetic and antihyperlipidemic remedies (Efferth and Kuete, 2010). Despite the presence of known antidiabetic medicine in the pharmaceutical market, diabetes and the related complications continued to be a major medical problem (Malviya et al., 2010). Antihyperglycemic effects of these plants are attributed to their ability to restore the function of pancreatic tissues by causing an increase in insulin output or inhibit the intestinal absorption of glucose or to the facilitation of metabolites in insulin dependent processes (Tiwari et al., 2014). However, More than 400 plant species having hypoglycemic activity have been available in literature, however, searching for new antidiabetic drugs from natural plants is still attractive because they contain substances which demonstrate alternative and safe effects on diabetes mellitus (Balamurugan et al., 2014). Most of plants contain glycosides, alkaloids, terpenoids, flavonoids, cartenoids, etc., that are frequently implicated as having antidiabetic effect. Species will be described in alphabetical order and information about each species will include in sequence: general botanical and taxonomic data, distribution in the world, experimental study and mechanism of action (Mngeni, 2017).

Traditional, complementary and alternative medicines have been used since ancient times (Harvey, 2008). Yet the use of traditional medicine (TM) remains widespread in developing countries, while use of complementary and alternative medicine (CAM) is increasing rapidly in developed countries. In many parts of the world, policy-makers, health professionals and the public are wrestling with questions about the safety, efficacy, quality, availability, preservation and further development of this type of health care (van Andel and Carvalheiro, 2013). TM is sometimes also the only affordable source of health care especially for the world's poorest patients (D'Amato et al., 2015).

Traditional medicine has been used for the treatment of human illnesses since long time and is mainly based on components derived from natural products, from herbs, plants, and animals. Medicinal natural products are very frequently used in Sudan and also are widely consumed in Africa and all over the world. About 80% of the populations in African countries depend on traditional medicine for their primary health care (Fabricant and Farnsworth, 2001). In Sudan, 90% of Sudan's population depends mainly on traditional medicine since admission to hospitals and obtaining modern synthetic drugs are limited and a high percentage of the population is nomads (Karar and Kuhnert, 2017). Sustainability of the use of medicinal plants is an important concern. The demand for medicinal plants is increasing in Africa as the population grows and pressure on medicinal plant resources will become greater than ever. Interest in plant-derived medicines has also increased in the developed countries among the pharmaceutical companies (Alves and Rosa, 2007). In contrast, due to their minor side effects, the medicinal plants are widely used to treat many human diseases (O'Hara et al., 1998).

A number of reviews from different countries have highlighted the significance of medicinal plants application for the control of diabetes (Upriety et al., 2012). In this review article, an attempt has been made to compile the reported hypoglycemic plants available in different scientific journals and may draw the attention useful to health professionals, scientists and scholars working in the field of pharmacology and therapeutics to develop evidence based alternative medicine to cure different kinds of diabetes in man and animals. However, this review shows the importance and the interest placed on Sudanese medicinal plants in the drive to demonstrate their antidiabetic effects and the responsible bioactive agents. This review also covers the common name of a plant, the parts that are commonly used as a remedy sources, extracts, doses, and a test model.

MATERIALS AND METHODS

Materials:

Publication regarding diabetes and effective plants were found in databases such as Science Direct, PubMed, Wiley, Scopus, and Springer. In addition to the above points of interest search Keywords used in this study included "medicinal plants", "diabetes", "Sudan", "herbal", and "treatment". Out of the 50 collected articles (published in the period between 1997 and 2018). As shown in (Table 1, 2 and 3).

Inclusion and exclusion criteria:

The search was restricted to English language articles. All studies found during the search were independently evaluated for competence and inclusion by two different authors. After compliance with inclusion criteria, experimental research and clinical trials that evaluate the effect of Sudanese medicinal herbs or plant component in diabetic animals or patients were included in the current research.

RESULTS AND DISCUSSION

Diabetes is a chronic disease that occurs when the body cannot produce enough insulin or cannot use insulin effectively (Bullon et al., 2014). It is projected that 300 million people will have the disease by the year 2025 (Mathers and Loncar, 2006) and it may reach to 366 million in the year 2030 (Hossain et al., 2009). Moreover, type 2 diabetes is a common condition and a serious global health problem (Narayan et al., 2000).

In most countries, diabetes has increased alongside rapid cultural and social changes: ageing populations, increasing urbanization, dietary changes, reduced physical activity and unhealthy behaviours (Petersen, 2003). Moreover, a person's risk of developing Type 2 Diabetes Mellitus has been shown to be highly linked to obesity and any family history or consequently of diabetes (Association, 2014).

Hyperglycaemic condition causes increased glycosylation leading to biochemical and morphological abnormalities due to altered protein structure and develop the neuropathy, retinopathy, neuropathy and cardiomyopathy (Kumar et al., 2011). All Statistical tabulation and analysis were done using Microsoft Excel Program (2016). Diabetes kills 1.1 million people in 2005 and more than 220 million people worldwide have diabetes, almost 80% of diabetes deaths occur in low- and middle-income countries (Cruz, 2007). The rising cost of medical care in Sudan is increasingly driving patients to herbal medicine and in case of diabetes, emerging experience with certain plants drew attention to some communities rarely known for such experience in the country.

Sudan is a developing country that frequently depends on folk medicine in all areas of the country. Several herbal preparations have been used in folklore practice for the management of diabetes with claims asserting their hypoglycemic effect. In this paper, an effort was made to refer to the different parts of 50 belonging to 32 families' plant species that are used in the Sudanese traditional medicine.

As shown in Table (1), detailed information about the plants and their antidiabetic activity from Sudan are presented. It was revealed that, up to 50 plant species belonging to 32 families were reported to have antidiabetic activity.

The family Fabaceae is considered a large family. The most common families are Fabaceae 9 species (18%) followed by Rubiaceae 6 species (12%), Amaryllidaceae, Apiaceae, Asteraceae, Poaceae and Zygophyllaceae 2 species (4%), Acanthaceae,

Apocynaceae, Arecaceae, Balanitaceae, Boraginaceae, Burseraceae, Capparaceae, Combretaceae, Cucurbitaceae, Cyperaceae, Ebenaceae, Euphorbiaceae, Lamiaceae, Lauraceae, Leguminosae, Martyniaceae, Meliaceae, Menispermaceae, Moringaceae, Myrtaceae, Pedaliaceae, Ranunculaceae, Rhamnaceae, Scrophulariaceae and Xanthorrhoeaceae one species (2%) shown in Table (2).

Distribution was prepared from family. The Fabaceae (*Lupinus albus*, *Trigonella foenum-graecum*, *Acacia albida*, *Acacia nilotica*, *Abrus precatorius*, *Bauhinia rufescens*, *Cicer arietinum* and *Acacia Senegal*). Rubiaceae (*Mitragyna inremis*, *Nauclea latifolia*, *Randia nilotica*, *Vangueria madagascariensis*, *Cephaelis ipecacuanha* and *Ruta graveolens*), Amaryllidaceae (*Allium Sativum* and *Allium cepa*), Apiaceae (*Foeniculum vulgare* and *Ammi visnaga*), Asteraceae (*Ambrosia maritima* and *Geigeria alata*), Poaceae (*Pennisetum glaucum* and *Cymbopogon proximus*) and Zygophyllaceae (*Zygophyllum coccineum* and *Balanites aegyptiaca*), Acanthaceae (*Blepharis liriifolia*), Apocynaceae (*Solenostemma argel*), Arecaceae (*Solenostemma argel*), Arecaceae (*Hyphaene thebaica*), Balanitaceae (*Balanites aegyptiaca*), Boraginaceae (*Cordia sinensis*), Burseraceae (*Boswellia papyrifera*), Capparaceae (*Capparis decidua*), Combretaceae (*Guiera senegalensis*), Cyperaceae (*Citrullus colocynthis*), Cyperaceae (*Cyperus rotundus*), Ebenaceae (*Diospyros mespiliformis*), Euphorbiaceae (*Croton zambiescus*), Lamiaceae (*Salvia officinalis*), Lauraceae (*Cinnamomum verum*), Leguminosae (*Lupinus albus*), Martyniaceae (*Martynia annua*), Meliaceae (*Khaya senegalensis*), Menispermaceae (*Tinospora bakis*), Moringaceae (*Moringa oleifera*), Myrtaceae (*Eucalyptus globulus*), Pedaliaceae (*Sesamum indicum*), Ranunculaceae (*Nigella sativa*), Rhamnaceae (*Ziziphus spina-christi*), Scrophulariaceae (*Striga hermonthica*) and Xanthorrhoeaceae (*Aloe sinkatana*).

The different parts plants and their antidiabetic Activity from Sudan are presented. Different plant parts including (Fruits, Leaves, Seeds, Roots, Whole plant, Stem bark, Bulb, Aerial parts, Epicarp, Grains, Mature Fruit, Pods, Rhizome, Root bark and Stem). There is a distinct preference for Fruits (23.08%), Leaves (17.31%), Seeds (15.38%), Roots (9.61%), Whole plant (7.69%), Stem bark (5.77%), Bulb (3.58%), Aerial parts (1.92%), Epicarp (1.92%), Grains (1.92%), Mature Fruit (1.92%), Pods (1.92%), Rhizome (1.92%), Root bark (1.92%) and Stem (1.92%) Table (2).

The Different used of solvent extracts (Aqueous, Ethanol, Methanol, Petroleum Ether, Dichloromethane and Ethyl Acetate). It was the highest percentage (%) of used solvent extracts Aqueous (48.61%), Ethanol (37.50%), Methanol (5.56%), Petroleum Ether, Dichloromethane and Ethyl Acetate (2.78%). The most used of extracts (95.74) and fractions (4.26%) Figure (1 & 2). Investigations of medicinal plants with different species and families were studied. Different parts of the plants were used for the antidiabetic study. The methanol, ethanol and aqueous solvents were most commonly used for the extractions. The preliminary phytochemical analyses mostly show the

presence of terpenoids and flavonoids. Efficacy evaluation of medicinal plants was done by streptozotocin or alloxan induced diabetic models. Most of the research results showed the hypoglycaemic effects and almost the same effect of standard drugs. Numerous mechanisms of action had been proposed for the plant extracts.

In addition, systematic and integrated studies on Sudanese medicinal plants in order to discover new antidiabetic drugs are absent; the majority of the studies are at the screening level on crude extracts.

The most commonly involved active constituents are Flavonoid, Tannin, Phenolics, and Alkaloid. Numerous mechanisms of actions have been proposed for these plant extracts. Some hypotheses relate to their effects on the activity of pancreatic β cells (synthesis, release) or the increase of the insulin sensitivity or the insulin-like activity of the plant extracts. All of these actions may be responsible for the reduction or abolition of diabetic complications.

CONCLUSION

Natural resources are still considered as potent candidates for drug discovery and are playing a pivotal role in drug development programs. Moreover, many medicinal herbs provide a rich mine for bioactive chemicals that are markedly free from undesirable side effects and of powerful pharmacological actions. The quest for control of diabetes has led to an increasing research at different fronts, among which is medicinal plants. Given the observation of an increasing use of medicinal plants for diabetes in Sudan, this necessitates validation of efficacy and safety. The attributed anti hyperglycaemic effects of these plants are due to their ability to restore the function of pancreatic tissues by causing an increase in insulin output or a decrease in the intestinal absorption of glucose. Hence, treatment with herbal drugs has effect on protecting cells and smoothing out fluctuation in glucose levels. In general, there is very little biological knowledge on the specific modes of action in the treatment of diabetes, but most of the plants have been found to contain substances like glycosides, alkaloids, terpenoids, flavonoids etc. that are frequently implicated as having anti diabetic effects. The research for alternative remedies (from the plant kingdom) for diabetes mellitus will continue all over the world as the disease poses many challenges not only to the physician but also to the researcher. However, data represented in this article prove a promising findings of getting a new herbal drugs which many reduce the heavy burden using synthetic compounds and hence wise the authors this article recommend more case studies involving standardized medicinal plant products should be carried

out in order to validate the usefulness of plant preparations in diabetes management, which will give support to the pre-clinical results. Consequently, much effort should be afforded to optimize a procedure for antidiabetic screening of different plants' extracts as well as isolated bioactive compounds for the discovery of new natural herbal antidiabetic drugs.

Table (1): Plants used in Sudanese traditional medicine for treatment of diabetes:

S/N	Plant Family	Plant Scientific Name	Plant Vernacular Name	Plant Part	Method used	Solvent used	Reference
1.	Acanthaceae	<i>Blepharis linariifolia</i>	Al-Bighail Shoak Al-Dhab	Whole plant	Extracts	Ethanol Aqueous	(Elbashir et al., 2018)
2.	Amaryllidaceae	<i>Allium Sativum</i>	Toom	Bulb	Extracts	Aqueous	(Kamal Eldin Gaber et al., 2013) (Eldin et al., 2009) (Ebrahim et al., 2012)
3.		<i>Allium cepa</i>	Basal	Bulb	Extract	Aqueous	(Gaber et al., 2013) (Eldin et al., 2009)
4.	Apiaceae	<i>Foeniculum vulgare</i>	El Shamar	Fruits	Extracts	Ethanolic	(Hilmi et al., 2014) (Yagi et al., 2013)
5.		<i>Ammi visnaga</i>	Khella	Fruits	Extracts	Ethanolic	(Hilmi et al., 2014) (Yagi et al., 2013)
6.	Apocynaceae	<i>Solenostemma argel</i>	Harjal	Leaves	Extracts	Ethanol Aqueous	(Elbashir et al., 2018)
7.	Arecaceae	<i>Hyphaene thebaica</i>	Nabag	Epicarp	Extracts	Aqueous	(Kamal Eldin Gaber et al., 2013) (Gaber et al., 2013)
8.	Asteraceae	<i>Ambrosia maritima</i>	Damesisa	Leaves	Extracts	Ethanolic	(Hilmi et al., 2014) (Yagi et al., 2013)
9.		<i>Geigeria alata</i>	El Gadgad	Aerial parts Roots	Extracts	Ethanol, Aqueous Methanol	(Elbashir et al., 2018) (Hafizur et al., 2012)
10.	Balanitaceae	<i>Balanites aegyptiaca</i>	Laloub	Fruit	Extracts	Aqueous	(Kamal Eldin Gaber et al., 2013)
11.	Boraginaceae	<i>Cordia sinensis</i>	Andrab	Leaves	Extracts	Ethanol Aqueous	(Elbashir et al., 2018)
12.	Burseraceae	<i>Boswellia papyrifera</i>	Tarag tarag, Shagar El-luban	Gum	Extracts	Ethanol Aqueous	(Elbashir et al., 2018)
13.	Capparaceae	<i>Capparis decidua</i>	Tundub	Stem	Extract	Methanol	(Zia-Ul-Haq et al., 2011)
14.	Combretaceae	<i>Guiera senegalensis</i>	Ghubeish	Leaves	Extracts	Ethanolic Aqueous	(Houacine et al., 2012) (Gaber et al., 2013) (Kamal Eldin Gaber et al., 2013)
15.	Cucurbitaceae	<i>Citrullus colocynthis</i>	Handal	Seeds	Extracts	Ethanol Aqueous	(Elbashir et al., 2018) (El Ghazali et al., 1997)
16.	Cyperaceae	<i>Cyperus rotundus</i>	Sieda	Rhizome Roots	Extracts	Ethanol Aqueous	(El Ghazali et al., 1997) (Elbashir et al., 2018)

17.	Ebenaceae	<i>Diospyros mespiliformis</i>	Gughan	stem barks	Fractio n	Petroleu m Ether Dichloro methane Ethyl Acetate Methanol	(Mohamed et al., 2009)
18.	Euphorbiaceae	<i>Croton zambiescus</i>	Um glela	Fruits	Fractio n	Petroleu m Ether Dichloro methane Ethyl Acetate Methanol	(Mohamed et al., 2009)
19.	Fabaceae	<i>Lupinus albus</i>	Tormus	Fruit	Extract s	Aqueous	(Kamal Eldin Gaber et al., 2013)
20.		<i>Trigonella foenum-graecum</i>	Hilba	Seed	Extract s	Aqueous	(Kamal Eldin Gaber et al., 2013)
21.		<i>Acacia albida</i>	Haraz	Root bark	Extract s	Aqueous	(Kamal Eldin Gaber et al., 2013)
22.		<i>Acacia nilotica</i>	Garad Sunt	Pods	Extract s	Ethanol Aqueous	(Elbashir et al., 2018)
23.		<i>Abrus precatorius</i>	Habbat Al-Arus	Seeds	Extract s	Ethanol Aqueous	(Elbashir et al., 2018)
24.		<i>Trigonella foenumgraceum</i>	Hilba	Seeds	Extract s	Ethanol Aqueous	(Elbashir et al., 2018) (Gaber et al., 2013)
25.		<i>Bauhinia rufescens</i>	Kulkul	Leaves	-	-	(El Ghazali et al., 1997)
26.		<i>Cicer arietinum</i>	Kabkabe	Seed	-	-	(Mustafa et al., 2013)
27.		<i>Acacia Senegal</i>	Hashab	Fruits	Extract s	Ethanolic	(Hilmi et al., 2014) (Yagi et al., 2013)
28.	Lamiaceae	<i>Salvia officinalis</i>	Meramia	Leaves	Extract s	Ethanolic	(Houacine et al., 2012)
29.	Lauraceae	<i>Cinnamomum verum</i>	Gerfa	Stem bark	Extract s	-	(Howeida et al., 2010)
30.	Leguminosae	<i>Lupinus albus</i>		-	Extract	Aqueous	(Gaber et al., 2013)
31.	Martyniaceae	<i>Martynia annua</i>	Gara Gebei	Mature Fruit	Extract s	Ethanol Aqueous	(Elbashir et al., 2018)
32.	Meliaceae	<i>Khaya senegalensis</i>	Mahogany	Stem bark	-	-	(El Ghazali et al., 1997)
33.	Menispermaceae	<i>Tinospora bakis</i>	Irg alhager	Seeds and Roots	Extract s	Ethanol Aqueous	(Alamin et al., 2015) (Elbashir et al., 2018)
34.	Moringaceae	<i>Moringa oleifera</i>	Moringa	Leaves	Extract s	Aqueous	(Selma et al., 2016)
35.	Myrtaceae	<i>Eucalyptus globulus</i>	ELBan; Kafour	Leaves	Extract s	Ethanolic	(Houacine et al., 2012)
36.	Pedaliaceae	<i>Sesamum indicum</i>	Simsem	Seeds	Extract s	Ethanolic	(Hilmi et al., 2014) (Yagi et al., 2013)
37.	Poaceae	<i>Pennisetum glaucum</i>	Dukhun	Grains	Extract s	Ethanol Aqueous	(Elbashir et al., 2018)
38.		<i>Cymbopogon proximus</i>	Mahareb	Whole plant	Extract s	Ethanol Aqueous	(Elbashir et al., 2018)

39.	Ranunculaceae	<i>Nigella sativa</i>	Kamon	Seeds	Extracts	Ethanol	(Hilmi et al., 2014) (Yagi et al., 2013)
40.	Rhamnaceae	<i>Ziziphus spina-christi</i>	Sidir	Roots	Extracts	Ethanol Aqueous	(Elbashir et al., 2018)
41.	Rubiaceae	<i>Mitragyna inremis</i>	Hikmat	Fruits	Extracts	Aqueous	(Alamin et al., 2015).
42.		<i>Nauclea latifolia</i>	Karmadoda	Fruits	Extracts	Aqueous	(Alamin et al., 2015)
43.		<i>Randia nilotica</i>	Kir Kir	Fruits	Extracts	Aqueous	(Alamin et al., 2015)
44.		<i>Vangueria madagascariensis</i>	Karkar	Fruit	Extracts	Ethanol Aqueous	(Elbashir et al., 2018)
45.		<i>Cephaelis ipecacuanha</i>	Irg Al-dahab	Roots	Extracts	Ethanol Aqueous	(Elbashir et al., 2018)
46.		<i>Ruta graveolens</i>	Sathab	Fruit	Extracts	Ethanol Aqueous	(Elbashir et al., 2018)
47.	Scrophulariaceae	<i>Striga hermonthica</i>	Boda	Whole plant	Extracts	Aqueous	(Alamin et al., 2015)
48.	Xanthorrhoeaceae	<i>Aloe sinkatana</i>	Sabar	Leaves	Extracts	Aqueous	(Kamal Eldin Gaber et al., 2013) (Gaber et al., 2013)
49.	Zygophyllaceae	<i>Zygophyllum coccineum</i>	Tartir	Whole plant	Extracts	Aqueous	(Kamal Eldin Gaber et al., 2013) (Gaber et al., 2013)
50.		<i>Balanites aegyptiaca</i>	Laloub	Fruit	Extract	Aqueous	(Gaber et al., 2013)

Table (2): Number and Percentage (%) of plant species tested for anti-diabetic activity peer plants family:

No.	Plant Family	Number of Plants	Percentage (%)
1.	Acanthaceae	1	2
2.	Amaryllidaceae	2	4
3.	Apiaceae	2	4
4.	Apocynaceae	1	2
5.	Arecaceae	1	2
6.	Asteraceae	2	4
7.	Balanitaceae	1	2
8.	Boraginaceae	1	2
9.	Burseraceae	1	2
10	Capparaceae	1	2
11	Combretaceae	1	2
12	Cucurbitaceae	1	2
13	Cyperaceae	1	2
14	Ebenaceae	1	2
15	Euphorbiaceae	1	2
16	Fabaceae	9	18
17	Lamiaceae	1	2
18	Lauraceae	1	2
19	Leguminosae	1	2
20	Martyniaceae	1	2
21	Meliaceae	1	2

22	Menispermaceae	1	2
23	Moringaceae	1	2
24	Myrtaceae	1	2
25	Pedaliaceae	1	2
26	Poaceae	2	4
27	Ranunculaceae	1	2
28	Rhamnaceae	1	2
29	Rubiaceae	6	12
30	Scrophulariaceae	1	2
31	Xanthorrhoeaceae	1	2
32	Zygophyllaceae	2	4
	Total	50	100

Table (2): Various parts of the plant are used and the number and Percentage (%) of plant species:

No.	Plant Parts Used	Number of Species	Percentage (%)
1.	Fruits	12	23.08
2.	Leaves	9	17.31
3.	Seeds	8	15.38
4.	Roots	5	9.61
5.	Whole plant	4	7.69
6.	Stem bark	3	5.77
7.	Bulb	2	3.85
8.	Aerial parts	1	1.92
9.	Epicarp	1	1.92
10.	Grains	1	1.92
11.	Gum	1	1.92
12.	Mature Fruit	1	1.92
13.	Pods	1	1.92
14.	Rhizome	1	1.92
15.	Root bark	1	1.92
16.	Stem	1	1.92
17.	Total	52	100

Figure (1): Percentage (%) of different used of extracts and Fractions.





Figure (2): Different uses of solvent to make plant extract.



ACKNOWLEDGEMENTS

We would like to thank group research of the International University of Africa (IUA), Faculty of Pure and Applied Sciences for financial support of the project. We also thank the leader of project **Prof. Dr.: Elbadri Elamin Osman**, Department of Microbiology, International University of Africa (IUA), Faculty of Pure and Applied Sciences, Khartoum, Sudan.

REFERENCES

Alamin, M. A., Yagi, A. I. & Yagi, S. M. 2015. Evaluation Of Antidiabetic Activity Of Plants Used In Western Sudan. *Asian Pacific Journal Of Tropical Biomedicine*, 5, 395-402.

Alves, R. R. & Rosa, I. M. 2007. Biodiversity, Traditional Medicine And Public Health: Where Do They Meet? *Journal Of Ethnobiology And Ethnomedicine*, 3, 14.

Anstee, Q. M., Targher, G. & Day, C. P. 2013. Progression Of Nafld To Diabetes Mellitus, Cardiovascular Disease Or Cirrhosis. *Nature Reviews Gastroenterology & Hepatology*, 10, 330.

Ashcroft, F. M. & Rorsman, P. 2012. Diabetes Mellitus And The B Cell: The Last Ten Years. *Cell*, 148, 1160-1171.

Association, A. D. 2014. Diagnosis And Classification Of Diabetes Mellitus. *Diabetes Care*, 37, S81-S90.

Association, A. D. 2015. 2. Classification And Diagnosis Of Diabetes. *Diabetes Care*, 38, S8-S16.

Association, A. D. 2016. Standards Of Medical Care In Diabetes—2016 Abridged For Primary Care Providers. *Clinical Diabetes: A Publication Of The American Diabetes Association*, 34, 3.

Balamurugan, K., Nishanthini, A. & Mohan, V. R. 2014. Antidiabetic And Antihyperlipidaemic Activity Of Ethanol

Extract Of *Melastoma Malabathricum* Linn. Leaf In Alloxan Induced Diabetic Rats. *Asian Pacific Journal Of Tropical Biomedicine*, 4, S442-S448.

Bullon, P., Newman, H. N. & Battino, M. 2014. Obesity, Diabetes Mellitus, Atherosclerosis And Chronic Periodontitis: A Shared Pathology Via Oxidative Stress And Mitochondrial Dysfunction? *Periodontology 2000*, 64, 139-153.

Cruz, A. A. 2007. *Global Surveillance, Prevention And Control Of Chronic Respiratory Diseases: A Comprehensive Approach*, World Health Organization.

D'amato, G., Holgate, S. T., Pawankar, R., Ledford, D. K., Cecchi, L., Al-Ahmad, M., Al-Enezi, F., Al-Muhsen, S., Ansotegui, I. & Baena-Cagnani, C. E. 2015. Meteorological Conditions, Climate Change, New Emerging Factors, And Asthma And Related Allergic Disorders. A Statement Of The World Allergy Organization. *World Allergy Organization Journal*, 8, 1.

Dunstan, D. W., Zimmet, P. Z., Welborn, T. A., De Courten, M. P., Cameron, A. J., Sicree, R. A., Dwyer, T., Colagiuri, S., Jolley, D. & Knuiman, M. 2002. The Rising Prevalence Of Diabetes And Impaired Glucose Tolerance: The Australian Diabetes, Obesity And Lifestyle Study. *Diabetes Care*, 25, 829-834.

Efferth, T. & Kuete, V. 2010. Cameroonian Medicinal Plants: Pharmacology And Derived Natural Products. *Frontiers In Pharmacology*, 1, 123.

El Ghazali, G., El Tohami, M. & El Egami, A. A. 1997. Medicinal Plants Of The Sudan. Part Iv. Medicinal Plants Of Northern Kordofan. *Khartoum: National Centre For Research Vii*.

Elbashir, S. M. I., Devkota, H. P., Wada, M., Kishimoto, N., Moriuchi, M., Shuto, T., Misumi, S., Kai, H. &

- & Watanabe, T. 2018. Free Radical Scavenging, A-Glucosidase Inhibitory And Lipase Inhibitory Activities Of Eighteen Sudanese Medicinal Plants. *Bmc Complementary And Alternative Medicine*, 18, 282.
- Eldin, I. M. T., Ahmed, E. M. & Elwahab, A. 2009. Clinical Hypoglycemic Effects Of Allium Cepa (Red Onion) In Type 1 Diabetic Patients. *Sudan Journal Of Medical Sciences*, 4.
- Fabricant, D. S. & Farnsworth, N. R. 2001. The Value Of Plants Used In Traditional Medicine For Drug Discovery. *Environmental Health Perspectives*, 109, 69.
- Gaber, K., Singhal, U. & Daowd, O. 2013. Hypoglycemic And Hypolipidaemic Effects Of Some Common Plants Extract In Type 2 Diabetic Patients At Eldabba Area (North Sudan). *Iosr J Pharm Biol Sci*, 8, 38-43.
- Hafizur, R. M., Babiker, R., Yagi, S., Chishti, S., Kabir, N. & Choudhary, M. I. 2012. The Antidiabetic Effect Of Geigeria Alata Is Mediated By Enhanced Insulin Secretion, Modulation Of B-Cell Function, And Improvement Of Antioxidant Activity In Streptozotocin-Induced Diabetic Rats. *Journal Of Endocrinology*, 214, 329-335.
- Harvey, A. L. 2008. Natural Products In Drug Discovery. *Drug Discovery Today*, 13, 894-901.
- He, Y., Yue, Y., Zheng, X., Zhang, K., Chen, S. & Du, Z. 2015. Curcumin, Inflammation, And Chronic Diseases: How Are They Linked? *Molecules*, 20, 9183-9213.
- Hilmi, Y., Abushama, M. F., Abdalgadir, H., Khalid, A. & Khalid, H. 2014. A Study Of Antioxidant Activity, Enzymatic Inhibition And In Vitro Toxicity Of Selected Traditional Sudanese Plants With Anti-Diabetic Potential. *Bmc Complementary And Alternative Medicine*, 14, 149.
- Hossain, P., Kavar, B. & El Nahas, M. 2009. Obesity And Diabetes In The Developing World—A Growing Challenge.
- Houacine, C., Elkhawad, A. O. & Ayoub, S. M. H. 2012. A Comparative Study On The Anti-Diabetic Activity Of Extracts Of Some Algerian And Sudanese Plants. *Journal Of Diabetes And Endocrinology*, 3, 25-28.
- Howeida, M. A., Idris, E. B., Almahdi, A. M., Sania, S. A., Abdelwahhab, M. H. & Mudawi, M. 2010. Antidiabetic And Hypolipidaemic Effects Of Cinnanomum Verum Bark On Hyperglycaemic And Diabetic Rats. *Research Journal Of Pharmacology*, 4, 21-25.
- Kahn, S. E., Hull, R. L. & Utzschneider, K. M. 2006. Mechanisms Linking Obesity To Insulin Resistance And Type 2 Diabetes. *Nature*, 444, 840.
- Kamal Eldin Gaber, P., Singhal, U. & Daowd, O. 2013. Hypoglycemic And Hypolipidaemic Effects Of Some Common Plants Extract In Type 2 Diabetic Patients At Eldabba Area (North Sudan).
- Karar, M. G. E. & Kuhnert, N. 2017. Herbal Drugs From Sudan: Traditional Uses And Phytoconstituents. *Pharmacognosy Reviews*, 11, 83.
- Kumar, D., Kumar, S., Kohli, S., Arya, R. & Gupta, J. 2011. Antidiabetic Activity Of Methanolic Bark Extract Of Albizia Odoratissima Benth. In Alloxan Induced Diabetic Albino Mice. *Asian Pacific Journal Of Tropical Medicine*, 4, 900-903.
- Malviya, N., Jain, S. & Malviya, S. 2010. Antidiabetic Potential Of Medicinal Plants. *Acta Pol Pharm*, 67, 113-118.
- Mathers, C. D. & Loncar, D. 2006. Projections Of Global Mortality And Burden Of Disease From 2002 To 2030. *Plos Medicine*, 3, E442.
- Matthews, D. & Matthews, P. 2011. Banting Memorial Lecture 2010. Type 2 Diabetes As An ‘Infectious’ Disease: Is This The Black Death Of The 21st Century? *Diabetic Medicine*, 28, 2-9.
- Mellitus, C. O. T. J. D. S. O. T. D. C. O. D., Seino, Y., Nanjo, K., Tajima, N., Kadowaki, T., Kashiwagi, A., Araki, E., Ito, C., Inagaki, N. & Iwamoto, Y. 2010. Report Of The Committee On The Classification And Diagnostic Criteria Of Diabetes Mellitus. *Journal Of Diabetes Investigation*, 1, 212-228.
- Mngeni, N. Z. 2017. *Bioactive Compounds From Selected Medicinal Plants Used In Antidiabetic Treatment*. Cape Peninsula University Of Technology.
- Mohamed, I. E., El Bushra, E., Choudhary, M. I. & Khan, S. N. 2009. Bioactive Natural Products From Two Sudanese Medicinal Plants Diospyros Mespiliformis And Croton Zambesicus. *Records Of Natural Products*, 3, 198.
- Mustafa, A., Eltayeb, B., Ali, M., Shaddad, A. & Mohammad, H. 2013. Antidiabetic And Hypolipidaemic Effects Of Cicer Arientinum Seedextracts In Hyperglycemic And Diabetic Rats. *J. Pharm. Biomed. Sci*, 30, 1046-1052.
- Narayan, K. V., Gregg, E. W., Fagot-Campagna, A., Engelgau, M. M. & Vinicor, F. 2000. Diabetes—A Common, Growing, Serious, Costly, And Potentially Preventable Public Health Problem. *Diabetes Research And Clinical Practice*, 50, S77-S84.
- O'hara, M., Kiefer, D., Farrell, K. & Kemper, K. 1998. A Review Of 12 Commonly Used Medicinal Herbs. *Archives Of Family Medicine*, 7, 523.
- Ogurtsova, K., Da Rocha Fernandes, J., Huang, Y., Linnenkamp, U., Guariguata, L., Cho, N., Cavan, D., Shaw, J. & Makaroff, L. 2017. Idf Diabetes Atlas: Global Estimates For The Prevalence Of Diabetes For 2015 And 2040. *Diabetes Research And Clinical Practice*, 128, 40-50.
- Patwardhan, B. & Partwardhan, A. 2005. *Traditional Medicine: Modern Approach For Affordable Global Health*, World Health Organization.
- Petersen, P. E. 2003. The World Oral Health Report 2003: Continuous Improvement Of Oral Health In The 21st Century—The Approach Of The Who Global Oral Health Programme. *Community Dentistry And Oral Epidemiology*, 31, 3-24.
- Selma, S., Khalid, H. & Ahmed, S. 2016. Anti-Diabetic Activity Of The Leaves Of Moringa Oleifera Lam. Growing In Sudan On Streptozotocin-Induced Diabetic Rats. *Brit. J. Med. Health Res*, 3, 48-57.
- Singab, A. N., Youssef, F. S. & Ashour, M. L. 2014. Medicinal Plants With Potential Antidiabetic Activity And Their Assessment. *Med Aromat Plants*, 3, 151.
- Singh, A. K. 2016. Exotic Ancient Plant Introductions: Part Of Indian ‘Ayurveda’ medicinal System. *Plant*

Genetic Resources, 14, 356-369.

Tiwari, N., Thakur, A., Kumar, V., Dey, A. & Kumar, V. 2014. Therapeutic Targets For Diabetes Mellitus: An Update. *Clinical Pharmacology & Biopharmaceutics*, 3, 1-10.

Upreti, Y., Asselin, H., Dhakal, A. & Julien, N. 2012. Traditional Use Of Medicinal Plants In The Boreal Forest Of Canada: Review And Perspectives. *Journal Of Ethnobiology And Ethnomedicine*, 8, 7.

Van Anandel, T. & Carvalheiro, L. G. 2013. Why Urban Citizens In Developing Countries Use Traditional Medicines: The Case Of Suriname. *Evidence-Based Complementary And Alternative Medicine*, 2013.

Yagi, S., Rahman, A. E. A., Elhassan, G. O. & Mohammed, A. M. 2013. Elemental Analysis Of Ten Sudanese Medicinal Plants Using X-Ray Fluorescence. *Journal Of Applied And Industrial Sciences*, 1, 49-53.

Yagi, S. M. & Yagi, A. I. 2018. Traditional Medicinal Plants Used For The Treatment Of Diabetes In The Sudan: A Review. *African Journal Of Pharmacy And Pharmacology*, 12, 27-40.

Zia-Ul-Haq, M., Cavar, S., Qayum, M., Imran, I. & Feo, V. D. 2011. Compositional Studies: Antioxidant And Antidiabetic Activities Of Capparis Decidua (Forsk.) Edgew. *International Journal Of Molecular Sciences*, 12, 8846-8861.